

# Incorporating area-level social drivers of health in predictive algorithms using electronic health record data

Agata Foryciarz<sup>1,\*</sup>, Nicole Gladish, PhD<sup>2</sup>, David H Rehkopf, ScD<sup>2,3,4,5,6</sup>, Sherri Rose, PhD<sup>3</sup>

<sup>1</sup>Department of Computer Science, Stanford University, Stanford, CA, USA

<sup>2</sup>Department of Epidemiology and Population Health, Stanford School of Medicine, Stanford, CA, USA

<sup>3</sup>Department of Health Policy, Stanford School of Medicine, Stanford, CA, USA

<sup>4</sup>Department of Medicine, Division of Primary Care and Population Health, Stanford School of Medicine, Stanford, CA, USA

<sup>5</sup>Department of Pediatrics, Stanford School of Medicine, Stanford, CA, USA

<sup>6</sup>Department of Sociology, Stanford University, Stanford, CA, USA

## **ABSTRACT**

The inclusion of social drivers of health (SDOH) into predictive algorithms of health outcomes has potential for improving algorithm performance, generalizability, transportability, and interpretation. However, there are limitations in the availability, understanding, and quality of SDOH variables, as well as a lack of guidance on how to incorporate them into algorithms when appropriate to do so. As such, few published algorithms include SDOH, and there is substantial methodological variability among those that do. We argue that practitioners should consider the use of social deprivation indices—a class of area-level SDOH measurements—given their accessibility, transparency, and quality. We illustrate the process of using such indices in predictive algorithms, which includes the selection of appropriate indices for the outcome, measurement time, and geographic level, in a demonstrative example with the Kidney Failure Risk Equation.

## INTRODUCTION

Social drivers of health (SDOH)—previously commonly referred to as social determinants of health—measure social structures and individual social factors that impact health, generate disease, and are among the most important contributors to health inequities.[1,2] SDOH encompass multiple domains, including economic stability, education access and quality, health care access and quality, neighborhood and built environment, and social and community context.[3]

There is potential that accounting for non-clinical factors impacting health can improve performance, generalizability, transportability, and interpretation of predictive algorithms of health outcomes. While some clinical algorithms contain SDOH, including cardiovascular risk prediction scores,[4,5] their incorporation in algorithms is not routine. This is due, in part, to a lack of recognition of the underlying processes that lead to observed health inequities. There are also problems in availability and quality of individual-level SDOH data.[6–8] It was only in 2024 that the Centers for Medicare and Medicaid Services started requiring SDOH screening for their hospital inpatient quality reporting program.[9] Published clinical algorithms that do use SDOH rely on heterogeneous data sources to calculate SDOH[10] and do not consistently provide detailed motivation for particular measurements. It is also the case that incorporating SDOH in predictive algorithms can reinforce health inequities.[11,12] Thus, it may not always be clear how to identify social factors relevant for a given clinical outcome, which SDOH data sources to use, or how to integrate them into algorithm development.

One category of well-studied and accessible SDOH variables is area based social deprivation indices, which describe social conditions in particular geographic areas.[13] In this article, we propose a set of considerations for incorporating social deprivation indices in predictive

algorithms using electronic health records data with the Kidney Failure Risk Equation (KFRE[14]) as an illustration. The KFRE is a risk stratification tool designed to identify chronic kidney disease (CKD) patients at highest risk of progression to kidney failure based on documented age, sex, estimated glomerular filtration rate (eGFR), and urine albumin creatinine ratio (uACR). It was developed using data from a single geographic region and external validation found a stable relationship between the predictors and outcome, attributing observed calibration differences to differences in baseline risk across study samples.[15] The inclusion of social factors in the predictive algorithm could help account for potential differences in baseline risk.

## **MOTIVATION FOR INCLUDING SOCIAL DRIVERS**

There is ample empirical evidence for associations between SDOH and health outcomes across disease areas.[16] While many health inequities stem from a common set of structural factors and are associated with mutually correlated SDOH,[16] different social drivers may have a more substantial contribution to particular health outcomes. Specific causal pathways may also vary across geographies and settings within them.[17] Additionally, domains of SDOH can be measured in many ways and at multiple levels—from individual to geographic area.[18,19] Hence, careful identification of SDOH domains and measurements relevant for a specific context before inclusion in a predictive algorithm is crucial. For the KFRE, factors across a range of domains, including those related to social status, stress, neighborhood and the health system, have been shown to play a role in CKD incidence and progression to kidney failure.[20–23] These factors contribute to existing disparities in the risk kidney failure incidence, which is 3.3 and 1.5 times higher for Black and Native Americans, compared to whites.[24,25]

Because SDOH are often strongly associated with health outcomes,[26] their inclusion in an algorithm may improve predictive performance. This has the potential to lead to improvements

in health equity, if, for example, more effective risk stratification leads to targeted interventions for at-risk populations. However, including SDOH in algorithms has relevance beyond improving predictive performance in the development cohort. When the development cohort represents a population with a heterogeneous distribution of SDOH associated with the outcome, including SDOH can improve algorithm generalizability to settings with a different distribution of SDOH.[27] Conversely, when the development cohort is homogenous with respect to SDOH, inclusion in the predictive algorithm may be less helpful. However, in these settings, assessing this SDOH homogeneity can still inform the feasibility of transportability to new populations.

Finally, consideration of SDOH is important from the perspective of having clear conceptual arguments for all predictive variables. In the absence of such conceptual clarity, imperfect proxies, such as race and ethnicity variables, are often used, reifying the erroneous use of race as a biological construct,[28] masking social processes,[29] and possibly contributing to racial health inequities by guiding care away from Black patients.[30]

### **AREA-LEVEL SOCIAL INDICES**

Social deprivation indices are composite area-level measurements often based on government data, such as the American Community Survey (ACS), that can be calculated at the state, county, census tract, census block group, or zip code level.[31] Sources of data for indices are scarce because only large-scale surveys are designed to be representative of small areas. Indices can represent multiple dimensions of deprivation with a single measure and have been associated with many health outcomes.[32–35]

Social deprivation indices are typically developed for identifying at-risk geographic areas to prioritize resource allocation. For example, the social vulnerability index (SVI) was developed for guiding natural disaster preparedness,[36] social deprivation index (SDI) for measuring health

care access,[32] index of concentration at the extremes (ICE) for measuring racial economic segregation,[33] and neighborhood stress score for risk adjustment.[37] They have also been applied in ways that extend beyond their original design. The SVI was used to guide COVID-19 testing and vaccine distribution[38] and the SDI is part of a cardiovascular risk score adapted by the American Heart Association.[5] Notably, most measures of social deprivation (with the exception of ICE) do not consider structural racism, which has a pronounced effect on health disparities.[33,39]

## **INCORPORATING INDICES IN PREDICTIVE ALGORITHMS**

Decisions about whether and how to include social deprivation indices among predictors will vary depending on the outcome, other selected predictors, and composition of the development cohort. These considerations are also relevant when choosing indices for study sample comparisons and stratified evaluation by index.

### **Identifying relevant indices with causal graphs**

The goal of including indices in a predictive algorithm is to capture relationships between social factors and the outcome. In the case of progression to kidney failure, these include access to care, economic and racial segregation, and neighborhood characteristics.[20] There is evidence of faster rates of CKD progression in Black and Native American populations[24,25] as well as delayed and lower-quality CKD care provided to Black patients.[23] Appropriate management of early to moderate CKD consists primarily of lifestyle counseling and pharmacological treatment as well as management of prevalent comorbid conditions.[40] We present a simplified representation of these possible causal processes in Figure 1A, distinguishing between area- and individual-level factors. Area income, segregation and neighborhood resources impact levels of access to healthcare, healthy food and safe physical activity, which in turn impact medical care received, diet, and physical activity.

Expressing these relationships with social deprivation indices involves identifying factors that incorporate unmeasured nodes in our causal graph. Two candidate indices fulfill those criteria: SDI and ICE (Figure 1B). SDI incorporates measures of poverty, nonemployment, household composition and housing quality, transportation and education.[32] ICE is a joint measure of racial and economic segregation.[33] Among its variants, we consider one which, for a given area  $i$  (with population size  $T_i$ ) compares the number of affluent white individuals (incomes  $\geq$  80th percentile nationally)  $A_i$  to the number of low-income non-white individuals ( $\leq$  20th percentile)  $P_i$ :  $ICE_i = (A_i - P_i)/T_i$ . While SDI and ICE measure overlapping concepts within SDOH domains (e.g., income percentile and percent living in poverty), they each were created to capture domains the other does not (e.g., racial segregation as a measure for structural racism, housing quality). As such, including both can be advantageous.

### **Geography and temporality**

Because indices can be calculated at various geographic levels, the same index might have different interpretations and associations with the outcome with consequences for generalizability and transportability.[19] Census tracts, for example, are more consistent in the number of people they capture than counties,[41] and zip codes are designed for delivering mail rather than capturing relatively homogenous geographic areas.[19,42] When neighboring geographic areas are heterogeneous with respect to a specific measure used to calculate an index (e.g., poverty), a larger geographic area that applies an average across them may mask those differences.[20] Meanwhile, other factors, such as measures of relative inequality within an area, may not be measurable in small areas. For our KFRE example, we use SDI and ICE indices at the census-tract level.

The time between measurement of variables used to calculate indices and the outcome may impact the validity of an index for a given predictive task, especially if indices show considerable variation over time. Prior empirical evidence can help inform how long it takes for factors captured in indices (e.g. income, food access, healthcare access) to affect a given outcome, and how rapidly the factors change over time. Because CKD develops slowly, exposures preceding kidney failure by as much as decades may be relevant for understanding current health. At the same time, current access to healthcare—captured by 2020 ACS-based indices—and effective management of comorbidities may have a larger effect on the speed of decline in later stages, which may be most relevant for the KFRE algorithm.

## **Distributions**

Examining the distributions of indices can help assess collinearity and generalizability. SDI and ICE are 81% correlated at the census-tract level in the general US population. We are interested in studying the KFRE in a primary care setting and introduce a US primary care cohort,[43] examining the SDI and ICE distributions in Figure 2. Figure 2A presents a joint distribution of the indices (correlation -0.83). While lower levels of SDI (lower deprivation) are associated with higher concentrations of wealthy, white individuals in a geographic area (and higher levels of SDI are associated with higher concentrations of poorer, non-white individuals), SDI values for areas with low levels of concentration at the extremes (near ICE=0) are spread across the entire range of the distribution. Figure 2B depicts the distributions of the two indices in the US population and the primary care cohort. Compared to the US population, the areas where individuals in the cohort reside tend to have higher concentrations of wealthy, white individuals and have higher levels of access to care, however, the populations overlap substantially.

## Use of indices for further evaluation of generalizability and transportability

Whether or not indices are included in the predictive algorithm, they can still be useful for reasoning about generalizability and transportability. This may be through stratified evaluation of algorithm performance across quantiles as well as comparison of the distribution of indices between development cohort and target populations or across different study samples. For example, if a distribution of an index differs noticeably between the development cohort and target population, this may suggest that additional validation is necessary to ensure that the relationship captured in the algorithm transports to the target population.[27]

## DISCUSSION

SDOH account for many health inequities and are important for designing appropriate interventions to reduce these inequities.[26,44–47] When clinical predictive algorithms are built with electronic health record data, the usefulness of the algorithm may be limited to individuals exposed to a similar, narrow set of social drivers. Consideration of relevant SDOH during algorithm development and evaluation can help validate algorithms and assess generalizability. Previously, no clear guidance has been available for identifying settings where incorporating social factors may be beneficial and how to do this rigorously.

**Table 1.** Guidance on considerations for inclusion of SDOH in predictive algorithms

Identifying an index relevant for the algorithm	<ul style="list-style-type: none"><li>● How has past literature related SDOH to the outcome?</li><li>● Collaborative creation of a causal graph describing underlying systems that created the data</li><li>● Which indices contribute to unmeasured nodes in the causal graph?</li><li>● How should indices be interpreted?</li><li>● Would including SDOH reinforce an unjust equilibrium?</li></ul>
Selecting an appropriate geographic level	<ul style="list-style-type: none"><li>● What levels are indices available for?</li><li>● Are concepts of interest more appropriately measured in larger or smaller areas?</li><li>● Do areas selected correspond to neighborhoods or other homogeneous environments?</li></ul>



Time of measurement	<ul style="list-style-type: none"> <li>• How long do factors captured in indices take to impact the outcome?</li> <li>• Does the index show considerable variability over time?</li> </ul>
Examining index distributions	<ul style="list-style-type: none"> <li>• How is the index distributed in the national population vs in the study sample?</li> <li>• If multiple indices are used, how correlated are they?</li> <li>• How to divide distribution into quantiles?</li> </ul>

We described such guidance for incorporating social deprivation indices in predictive algorithms (summarized in Table 1) and the implications for performance, generalizability, transportability, and interpretation. These indices have several advantages, including their availability and validation. Despite this, they may not be the most appropriate SDOH variables to include if they do not sufficiently capture causal paths or reinforce unjust equilibriums in the healthcare system. Indices reflect conditions of an individual’s environment, which may differ from individual-level factors, and when individual-level factors are most relevant, indices may not be appropriate proxies.[48,49] Moreover, while they have been used in modeling health outcomes,[4,5] the indices were developed primarily for purposes other than building predictive algorithms. Additional validation is warranted for ascertaining whether they are appropriate to use in specific contexts. Finally, the majority of measures of social deprivation (with the exception of ICE, which captures racial-economic segregation) do not consider structural racism, unlike other measures.[50,51]

**FUNDING**

AF and SR were supported by NIH grant R01LM013989. NG and DHR were supported by Robert Wood Johnson Foundation Grant #79319.

**AUTHOR CONTRIBUTIONS**

All authors contributed to project conceptualization and formative discussions. AF analyzed PRIME data, under supervision of SR, and wrote the original draft. NG generated and curated social deprivation indices data, in collaboration with DHR. All authors reviewed and edited the manuscript.

## **ACKNOWLEDGEMENTS**

The authors thank Professor Alyce Adams, Dr. Robert Phillips, and Dr. Andrew Bazemore for their insights on this work. The authors also thank Malcolm Barrett for code review.

## **CONFLICT OF INTEREST STATEMENT**


None declared.

## **DATA AVAILABILITY**

The following acknowledgment text is included regarding data availability as described by the Stanford Center for Population Health Sciences Data Core (<https://phsdocs.stanford.edu/v1.0/need-help/citing-phs-data-core>): “Data for this project were accessed using the Stanford Center for Population Health Sciences Data Core. The PHS Data Core is supported by a National Institutes of Health National Center for Advancing Translational Science Clinical and Translational Science Award (UL1TR003142) and from Internal Stanford funding. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.”


## FIGURE LEGENDS

**Figure 1.** Possible causal graphs representing the relationships between a selected subset of SDOH and variables used in KFRE. Measured variables are shown in yellow (if they are part of KFRE) or orange (if they are not). Unmeasured variables are shown in blue. A. Causal graph with SDOH variables. B. Causal graph with a subset of SDOH variables represented by selected indices (SDI, ICE) in red.




**Figure 2.** Distributions of 2020 ICE and SDI in a US primary care cohort at the census-tract level. SDI ranges from 0 to 100, with 100 indicating higher levels of deprivation. ICE ranges from -1 to 1, with -1 indicating higher levels of inequality. A. Joint distribution of the two indices. B. Comparisons of the index distributions in the cohort (red) and the general US population (blue) with mean values denoted using vertical lines. The mean values in part B are weighed by the population of each census tract (for US) or number of people in the given census tract (Cohort). In all figures, bars and cells corresponding to fewer than 11 individuals were suppressed for data privacy.

A.



B.



## REFERENCES

- 1 Marmot M, Friel S, Bell R, et al. Closing the gap in a generation: health equity through action on the social determinants of health. *Lancet*. 2008;372:1661–9.
- 2 World Health Organization. Social Determinants of Health. World Health Organization. [www.who.int/health-topics/social-determinants-of-health](http://www.who.int/health-topics/social-determinants-of-health) (accessed 14 January 2024)
- 3 Pronk N, Kleinman DV, Goekler SF, et al. Promoting Health and Well-being in Healthy People 2030. *J Public Health Manag Pract*. 2021;27:S242–8.
- 4 Hippisley-Cox J, Coupland C, Vinogradova Y, et al. Derivation and validation of QRISK, a new cardiovascular disease risk score for the United Kingdom: prospective open cohort study. *BMJ*. 2007;335:136.
- 5 Khan SS, Coresh J, Pencina MJ, et al. Novel Prediction Equations for Absolute Risk Assessment of Total Cardiovascular Disease Incorporating Cardiovascular-Kidney-Metabolic Health: A Scientific Statement From the American Heart Association. *Circulation*. 2023;148:1982–2004.
- 6 Wang M, Pantell MS, Gottlieb LM, et al. Documentation and review of social determinants of health data in the EHR: measures and associated insights. *J Am Med Inform Assoc*. 2021;28:2608–16.
- 7 Cook LA, Sachs J, Weiskopf NG. The quality of social determinants data in the electronic health record: a systematic review. *J Am Med Inform Assoc*. 2021;29:187–96.
- 8 Linfield GH, Patel S, Ko HJ, et al. Evaluating the comparability of patient-level social risk data extracted from electronic health records: A systematic scoping review. *Health Informatics J*. 2023;29:14604582231200300.
- 9 Department of Health and Human Services, Centers For Medicare & Medicaid Services. Medicare Program; Hospital Inpatient Prospective Payment Systems for Acute Care Hospitals and the Long-Term Care Hospital Prospective Payment System and Policy Changes and Fiscal Year 2023 Rates; Quality Programs and Medicare Promoting Interoperability Program Requirements for Eligible Hospitals and Critical Access Hospitals; Costs Incurred for Qualified and Non-Qualified Deferred Compensation Plans; and Changes to Hospital and Critical Access Hospital Conditions of Participation. Final Rule. 2022;87:48780–9499.
- 10 Chen M, Tan X, Padman R. Social determinants of health in electronic health records and their impact on analysis and risk prediction: A systematic review. *J Am Med Inform Assoc*. 2020;27:1764–73.
- 11 McWilliams JM, Weinreb G, Ding L, et al. Risk adjustment and promoting health equity in population-based payment: Concepts and evidence. *Health Aff (Millwood)*. 2023;42:105–14.

- 12 Bergquist SL, Layton TJ, McGuire TG, et al. Data transformations to improve the performance of health plan payment methods. *J Health Econ.* 2019;66:195–207.
- 13 Gladish N, Phillips R, Ouyang D, Hao S, Chu I, Rehkopf D. Social Deprivation and Vulnerability Indices: SVI, ADI, SDI, NSS7, FDep, ICE. 2022. <https://doi.org/10.57761/75cc-1t35>
- 14 Tangri N, Stevens LA, Griffith J, et al. A predictive model for progression of chronic kidney disease to kidney failure. *JAMA.* 2011;305:1553–9.
- 15 Tangri N, Grams ME, Levey AS, et al. Multinational Assessment of Accuracy of Equations for Predicting Risk of Kidney Failure: A Meta-analysis. *JAMA.* 2016;315:164–74.
- 16 Solar O, Irwin A. A conceptual framework for action on the social determinants of health. Published Online First: 2010.
- 17 Kunitz SJ. *The Health of Populations: General Theories and Particular Realities.* Oxford University Press 2007.
- 18 Braveman PA, Cubbin C, Egerter S, et al. Socioeconomic status in health research: one size does not fit all. *JAMA.* 2005;294:2879–88.
- 19 Krieger N, Chen JT, Waterman PD, et al. Geocoding and monitoring of US socioeconomic inequalities in mortality and cancer incidence: does the choice of area-based measure and geographic level matter?: the Public Health Disparities Geocoding Project. *Am J Epidemiol.* 2002;156:471–82.
- 20 Norton JM, Moxey-Mims MM, Eggers PW, et al. Social Determinants of Racial Disparities in CKD. *J Am Soc Nephrol.* 2016;27:2576–95.
- 21 Hannan M, Ansari S, Meza N, et al. Risk Factors for CKD Progression: Overview of Findings from the CRIC Study. *Clin J Am Soc Nephrol.* 2021;16:648–59.
- 22 Crews DC, Liu Y, Boulware LE. Disparities in the burden, outcomes, and care of chronic kidney disease. *Curr Opin Nephrol Hypertens.* 2014;23:298–305.
- 23 Eneanya ND, Boulware LE, Tsai J, et al. Health inequities and the inappropriate use of race in nephrology. *Nat Rev Nephrol.* 2022;18:84–94.
- 24 McClellan W, Warnock DG, McClure L, et al. Racial Differences in the Prevalence of Chronic Kidney Disease among Participants in the Reasons for Geographic and Racial Differences in Stroke (REGARDS) Cohort Study. *J Am Soc Nephrol.* 2006;17:1710.
- 25 Saran R, Robinson B, Abbott KC, et al. US Renal Data System 2016 Annual Data Report: Epidemiology of Kidney Disease in the United States. *Am J Kidney Dis.* 2017;69:A7–8.
- 26 Hood CM, Gennuso KP, Swain GR, et al. County Health Rankings: Relationships Between Determinant Factors and Health Outcomes. *Am J Prev Med.* 2016;50:129–35.
- 27 Degtiar I, Rose S. A Review of Generalizability and Transportability. *Annual Review of Statistics and Its Application.* 2023;10:501–24.

- 28 Cerdeña JP, Plaisime MV, Tsai J. From race-based to race-conscious medicine: how anti-racist uprisings call us to act. *Lancet*. 2020;396:1125–8.
- 29 Chowkwanyun M. THE STRANGE DISAPPEARANCE OF HISTORY FROM RACIAL HEALTH DISPARITIES RESEARCH. *Du Bois Rev*. 2011;8:253–70.
- 30 Vyas DA, Eisenstein LG, Jones DS. Hidden in Plain Sight — Reconsidering the Use of Race Correction in Clinical Algorithms. *N Engl J Med*. 2020;383:874–82.
- 31 Rehkopf DH, Glymour MM, Osypuk TL. The Consistency Assumption for Causal Inference in Social Epidemiology: When a Rose is Not a Rose. *Curr Epidemiol Rep*. 2016;3:63–71.
- 32 Butler DC, Petterson S, Phillips RL, et al. Measures of social deprivation that predict health care access and need within a rational area of primary care service delivery. *Health Serv Res*. 2013;48:539–59.
- 33 Krieger N, Waterman PD, Spasojevic J, et al. Public Health Monitoring of Privilege and Deprivation With the Index of Concentration at the Extremes. *Am J Public Health*. 2016;106:256–63.
- 34 Bevan GH, Nasir K, Rajagopalan S, et al. Socioeconomic Deprivation and Premature Cardiovascular Mortality in the United States. *Mayo Clin Proc*. 2022;97:1108–13.
- 35 Kind AJH, Jencks S, Brock J, et al. Neighborhood socioeconomic disadvantage and 30-day rehospitalization: a retrospective cohort study. *Ann Intern Med*. 2014;161:765–74.
- 36 Flanagan BE, Gregory EW, Hallisey EJ, et al. A Social Vulnerability Index for Disaster Management. *Journal of Homeland Security and Emergency Management*. 2011;8. doi: 10.2202/1547-7355.1792
- 37 Ash AS, Mick EO, Ellis RP, et al. Social Determinants of Health in Managed Care Payment Formulas. *JAMA Intern Med*. 2017;177:1424–30.
- 38 Center for Disease Control. Innovative Uses of SVI During COVID-19. Agency for Toxic Substances and Disease Registry. [https://www.atsdr.cdc.gov/placeandhealth/project\\_snapshots/svitool\\_covid.html](https://www.atsdr.cdc.gov/placeandhealth/project_snapshots/svitool_covid.html) (accessed 15 January 2024)
- 39 Yearby R, Clark B, Figueroa JF. Structural Racism In Historical And Modern US Health Care Policy. *Health Aff*. 2022;41:187–94.
- 40 Wouters OJ, O'Donoghue DJ, Ritchie J, et al. Early chronic kidney disease: diagnosis, management and models of care. *Nat Rev Nephrol*. 2015;11:491–502.
- 41 United States. Bureau of the Census. Geographic Areas Reference Manual. U.S. Department of Commerce, Economics and Statistics Administration, Bureau of the Census 1994.
- 42 Grubesic TH, Matisziw TC. On the use of ZIP codes and ZIP code tabulation areas (ZCTAs) for the spatial analysis of epidemiological data. *Int J Health Geogr*. 2006;5:58.
- 43 Phillips R. The PRIME registry helps thousands of primary care clinicians liberate EHR data and prepare for MIPS. Published Online First: 2017.

- 44 Thornton RLJ, Glover CM, Cené CW, et al. Evaluating Strategies For Reducing Health Disparities By Addressing The Social Determinants Of Health. *Health Aff* . 2016;35:1416–23.
- 45 Haas JS, Linder JA, Park ER, et al. Proactive tobacco cessation outreach to smokers of low socioeconomic status: a randomized clinical trial. *JAMA Intern Med*. 2015;175:218–26.
- 46 Bloch G, Rozmovits L. Implementing social interventions in primary care. *CMAJ*. 2021;193:E1696–701.
- 47 Gottlieb LM, Wing H, Adler NE. A Systematic Review of Interventions on Patients' Social and Economic Needs. *Am J Prev Med*. 2017;53:719–29.
- 48 Moss JL, Johnson NJ, Yu M, et al. Comparisons of individual- and area-level socioeconomic status as proxies for individual-level measures: evidence from the Mortality Disparities in American Communities study. *Popul Health Metr*. 2021;19:1.
- 49 Brown EM, Franklin SM, Ryan JL, et al. Assessing Area-Level Deprivation as a Proxy for Individual-Level Social Risks. *Am J Prev Med*. 2023;65:1163–71.
- 50 Hardeman RR, Homan PA, Chantarat T, et al. Improving the measurement of structural racism to achieve antiracist health policy. *Health Aff (Millwood)*. 2022;41:179–86.
- 51 Groos M, Wallace M, Hardeman R, et al. Measuring inequity: a systematic review of methods used to quantify structural racism. *J Health Dispar Res Pract*. 2018;11:13.